

QUANTIZATION OF MINIMAL RESOLUTIONS OF KLEINIAN SINGULARITIES

MITYA BOYARCHENKO

ABSTRACT. In this paper we prove an analogue of a recent result of Gordon and Stafford that relates the representation theory of certain noncommutative deformations of the coordinate ring of the n -th symmetric power of $\mathbb{C}^2$ with the geometry of the Hilbert scheme of n points in $\mathbb{C}^2$ through the formalism of $\mathbb{Z}$ -algebras. Our work produces, for every *regular* noncommutative deformation $\mathcal{O}^\lambda$ of a Kleinian singularity $X = \mathbb{C}^2/\Gamma$, as defined by Crawley-Boevey and Holland, a filtered $\mathbb{Z}$ -algebra which is Morita equivalent to $\mathcal{O}^\lambda$, such that the associated graded $\mathbb{Z}$ -algebra is Morita equivalent to the minimal resolution of X . The construction uses the description of the algebras $\mathcal{O}^\lambda$ as quantum Hamiltonian reductions, due to Holland, and a GIT construction of minimal resolutions of X , due to Cassens and Slodowy.

CONTENTS

1. Introduction	1
2. A study of the minimal resolution	5
3. Quantization of Kleinian singularities	9
4. Morita $\mathbb{Z}$ -algebras	11
5. Quantization of the minimal resolution	16
6. Proof of the main theorem	18
References	24

1. INTRODUCTION

1.1. Let $\Gamma \subset SL_2(\mathbb{C})$ be a finite subgroup, and $\{\mathcal{O}^\lambda\}$ the family of noncommutative deformations of the singularity $X = \mathbb{C}^2/\Gamma$ constructed by Crawley-Boevey and Holland [CBH]. If (Q, I) denotes the McKay quiver associated to Γ , then the parameter space for these deformations is naturally identified with $\mathbb{C}^I$. Explicitly, given $\lambda \in \mathbb{C}^I$, recall ([CBH], p. 606) that the *deformed preprojective algebra* of Q with parameter λ is defined by

$$\Pi^\lambda = \Pi^\lambda(Q) = \mathbb{C}\overline{Q} \Big/ \left\langle \left\langle \sum_{a \in Q} [a, a^*] - \lambda \right\rangle \right\rangle,$$

The author's research was partially supported by NSF grant DMS-0401164.
Address: Department of Mathematics, University of Chicago, Chicago, IL 60637
E-mail: mitya@math.uchicago.edu.

and $\mathcal{O}^\lambda$ is the spherical subalgebra $\mathcal{O}^\lambda = e_0 \Pi^\lambda e_0$. Here $\overline{Q}$ denotes the *double of Q* , i.e., the quiver obtained from Q by adding an arrow a^* for each arrow $a \in Q$ such that the tail (resp., head) of a^* is the head (resp., tail) of a . We write $\mathbb{C}\overline{Q}$ for the path algebra of $\overline{Q}$, and $e_i \in \mathbb{C}\overline{Q}$ for the idempotent corresponding to the vertex $i \in I$; the extending vertex of Q is denoted by $0 \in I$. Finally, λ is identified with the element $\sum_{i \in I} \lambda_i e_i \in \mathbb{C}\overline{Q}$.

We define a grading on the algebra $\mathbb{C}\overline{Q}$ by assigning degree 0 to each idempotent e_i , and degree 1 to each arrow $a \in Q$ and its opposite arrow a^* ; in other words, the grading is by the length of paths. The algebras Π^λ and $\mathcal{O}^\lambda$ inherit natural filtrations from this grading. This filtration will be used throughout the paper without explicit mention.

1.2. We let $\delta \in \mathbb{N}^I$ denote the minimal positive imaginary root for Q , and we write

$$\begin{aligned} \Lambda &= \{\xi \in \mathbb{Z}^I \mid \xi \cdot \delta = 0\}, \\ \Lambda_+ &= \{\xi \in \Lambda \mid \xi \cdot \alpha \geq 0 \text{ for every positive Dynkin root } \alpha\}, \\ \Lambda_{++} &= \{\xi \in \Lambda \mid \xi \cdot \alpha > 0 \text{ for every positive Dynkin root } \alpha\}. \end{aligned}$$

Hereafter, a *root* is an element of the root system associated to the quiver Q , which in this case can be defined as the set of all $\alpha \in \mathbb{Z}^I \setminus \{0\}$ such that $q(\alpha) \leq 1$, where q is the Tits form corresponding to Q . A root α is *Dynkin* if $\alpha \cdot \epsilon_0 = 0$, where $\epsilon_0 \in \mathbb{Z}^I$ is the standard coordinate vector corresponding to the extending vertex (in other words, the coordinate of α corresponding to the extending vertex is zero). A root α is *real* (resp., *imaginary*) if $q(\alpha) = 1$ (resp., $q(\alpha) = 0$); note that a Dynkin root is automatically real. There is a natural identification of Λ with the weight lattice of the finite root system associated to the Dynkin diagram obtained by deleting the extending vertex, so that Λ_+ (resp., Λ_{++}) corresponds to the set of dominant (resp., dominant regular) weights.

1.3. We denote by $\text{Rep}(Q, \delta)$ (resp., $\text{Rep}(\overline{Q}, \delta)$) the affine space of all representations of Q (resp., $\overline{Q}$) with dimension vector δ . Using the trace pairing, $\text{Rep}(\overline{Q}, \delta)$ is naturally identified with the cotangent bundle $T^* \text{Rep}(Q, \delta)$. Let

$$G = PGL(\delta) = \left(\prod_{i \in I} GL(\delta_i, \mathbb{C}) \right) / \mathbb{C}^\times,$$

where $\mathbb{C}^\times$ is embedded diagonally into the product. This is a reductive algebraic group acting by conjugation on the varieties $\text{Rep}(Q, \delta)$ and $\text{Rep}(\overline{Q}, \delta)$, and we write

$$\mu : \text{Rep}(\overline{Q}, \delta) \longrightarrow \mathfrak{g}^*$$

for the moment map for the action of G on $\text{Rep}(Q, \delta)$ (see [CBH], p. 606), where

$$\mathfrak{g} = \text{Lie}(G) = \mathfrak{p}\mathfrak{gl}(\delta) = \left(\prod_{i \in I} \mathfrak{gl}(\delta_i, \mathbb{C}) \right) / \mathbb{C}$$

is the Lie algebra of G .

1.4. Using the determinant maps $\det : GL(\delta_i, \mathbb{C}) \rightarrow \mathbb{C}^\times$, we identify Λ with the group of 1-dimensional characters of G . Given $\chi \in \Lambda_{++}$, Cassens and Slodowy [CS] construct a minimal resolution of the Kleinian singularity X as the projective morphism

$$\tilde{X} := \text{Proj } S \longrightarrow \text{Spec } S_0 \cong X,$$

where S is the graded algebra

$$S = \bigoplus_{n \geq 0} S_n, \quad S_n = \mathbb{C}[\mu^{-1}(0)]^{G, \chi^n}. \quad (1.1)$$

Here $\mathbb{C}[\mu^{-1}(0)]$ stands for the algebra of regular functions on the *scheme-theoretic* fiber of μ over $0 \in \mathfrak{g}^*$ and $\mathbb{C}[\mu^{-1}(0)]^{G, \chi^n}$ denotes the G -eigenspace corresponding to the character χ^n . Note that each component S_n of S is itself graded, where the grading is induced by the grading on $\mathbb{C}[\mu^{-1}(0)]$, which in turn is induced by the grading of $\mathbb{C}[\text{Rep}(\overline{Q}, \delta)]$ by the degree of polynomials (we use the fact that $\text{Rep}(\overline{Q}, \delta)$ is an affine space).

The minimal resolution $\tilde{X} \rightarrow X$ is studied in more detail in Section 2, where we also prove a result (Theorem 2.2) on the structure of the ring S that, to the best of our knowledge, does not appear in the existing literature.

1.5. Let $\lambda \in \mathbb{C}^I$ be such that $\lambda \cdot \delta = 1$ and $\mathcal{O}^\lambda$ is regular, i.e., has finite global dimension. By Theorem 0.4 in [CBH], regularity in this case is equivalent to either of the following two properties: $\mathcal{O}^\lambda$ is Morita equivalent to the deformed preprojective algebra Π^λ , or $\lambda \cdot \alpha \neq 0$ for every Dynkin root α . Using Holland's description [Ho] of the algebras $\mathcal{O}^\lambda$ as quantum Hamiltonian reductions of the algebra of polynomial differential operators on $\text{Rep}(Q, \delta)$, which is recalled in Section 3, we construct, for a given $\chi \in \Lambda_{++}$, a filtered $\mathbb{Z}$ -algebra $B^\lambda(\chi)$ such that

- (1) $\mathcal{O}^\lambda\text{-mod}$ is naturally equivalent to $B^\lambda(\chi)\text{-qgr}$, in a way compatible with filtrations, and
- (2) there is a natural isomorphism $\text{gr}^\bullet B^\lambda(\chi) \cong \hat{S}$, where $\hat{S}$ is the $\mathbb{Z}$ -algebra associated to the graded algebra S defined by (1.1).

The construction is described in Section 5, where we also state our main result, Theorem 5.2, which is then proved in Section 6.

1.6. The $\mathbb{Z}$ -algebra $B^\lambda(\chi)$ has the property that $B^\lambda(\chi)_{0,0} = \mathcal{O}^{\lambda+\xi}$ for a “sufficiently large” $\xi \in \Lambda_{++}$, and then

$$B^\lambda(\chi)_{n,n} = \mathcal{O}^{\lambda+n \cdot \chi + \xi} \quad \text{for all } n \geq 0.$$

Thus, the construction of $B^\lambda(\chi)$ depends on the choice of ξ . However, this dependence is not very serious, since two different choices of ξ lead to naturally Morita equivalent $\mathbb{Z}$ -algebras, which is why ξ is omitted from the notation. Furthermore, if $\mathcal{O}^\lambda$ has no nonzero finite dimensional modules, one can take $\xi = 0$, and we conjecture that one can always take $\xi = 0$ as long as λ is *dominant* (see Remark 6.10).

More importantly, $B^\lambda(\chi)$ also depends on the choice of $\chi \in \Lambda_{++}$. For this reason our quantization of the minimal resolution $\tilde{X}$ may be called “non-canonical.” A more canonical version of the quantization would consist of replacing $B^\lambda(\chi)$ by a “lower-triangular Λ -algebra,” bigraded by Λ_+ instead of $\mathbb{Z}_+$, and realizing $\tilde{X}$ as a suitable “multi- Proj ” of the Λ_+ -graded ring $\bigoplus_{\chi \in \Lambda_+} \mathbb{C}[\mu^{-1}(0)]^{G, \chi}$. Note, however, that such a construction cannot be obtained by a straightforward modification of the results of the present paper. The most apparent reason for this is that we repeatedly make crucial use of the following simple fact: given a natural number N , every integer $n \geq 2N - 1$ can be written as a sum of integers that lie between N and $2N - 1$. However, this fact has no suitable analogue for lattices other than $\mathbb{Z}$, in the sense that if $\text{rk } \Lambda \geq 2$, then Λ_{++} is *not* finitely generated as a monoid. This issue will be addressed in [B].

1.7. A canonical (in the sense explained above) quantization of minimal resolutions of Kleinian singularities of type A has been recently constructed by I. Musson [Mu]. His approach is very different from ours in that instead of using Holland’s results, he constructs a filtered $\mathbb{Z}$ -algebra that deforms the minimal resolution by using the explicit description of the latter as a toric variety (which replaces Cassens and Slodowy’s construction). In particular, this approach does not generalize to other types of Kleinian singularities. Apart from the basic theory of $\mathbb{Z}$ -algebras, our papers are completely disjoint, and can be read independently.

1.8. We end the introduction by collecting some of the notation and terminology that is used throughout the paper. A ring is said to be *regular* if it has finite global homological dimension. If A is a ring, then $A\text{-Mod}$ and $A\text{-mod}$ denote the category of left (resp., left finitely generated) A -modules. More generally, we follow the conventions of [GS] and [Mu], in that any notation involving an uppercase letter denotes a category of all modules with a certain property, whereas the corresponding notation involving only lowercase letters denotes the full subcategory consisting of finitely generated modules.

In particular, if $A = \bigoplus_{n \geq 0} A_n$ is a graded algebra, we denote by $A\text{-Grmod}$ the category of $\mathbb{Z}$ -graded (not necessarily positively graded) A -modules. A graded A -module M is said to be *bounded* if $M_n = (0)$ for $n \gg 0$, and we denote by $A\text{-Tors} \subseteq A\text{-Grmod}$ the full subcategory consisting of modules that are unions of their bounded submodules. The corresponding Serre quotient

$$A\text{-Qgr} = \frac{A\text{-Grmod}}{A\text{-Tors}}$$

plays the role of the “category of quasi-coherent sheaves on $\text{Proj } A$.” According to the previous paragraph, we have the corresponding abelian categories $A\text{-grmod}$, $A\text{-tors}$ and $A\text{-qgr}$. The notion of a (lower-triangular) $\mathbb{Z}$ -algebra B , as well as the associated abelian categories $B\text{-Qgr}$, $B\text{-qgr}$, etc., are introduced in Section 4.

An *algebra* will always mean for us an algebra over $\mathbb{C}$, and if A, B are algebras, then an (A, B) -bimodule M is required to satisfy the condition that the two induced actions of $\mathbb{C}$ on M coincide. All tensor products, unless specified otherwise, will be taken over $\mathbb{C}$. With

the exception of $\mathbb{Z}$ -algebras, all rings are assumed to have a multiplicative identity, and all modules are assumed to be unital.

1.9. Acknowledgements. I am greatly indebted to Victor Ginzburg for introducing me to this area of research, stating the problem, and constant encouragement and attention to my work. During our numerous conversations he suggested several ideas that were crucial for some of the proofs appearing in this paper. I am also grateful to Dennis Gaitsgory for motivating discussions during the early stages of this work. Finally, I would like to thank Ian Musson for explaining his work to me and sharing his preprint [Mu] before it was made available to the general audience.

I am thankful to Victor Ginzburg, Iain Gordon and Ian Musson for making helpful comments, suggesting improvements, and pointing out several misprints in the first version of the paper.

2. A STUDY OF THE MINIMAL RESOLUTION

2.1. Recall the notation $X = \mathbb{C}^2/\Gamma$ and $\mu : \text{Rep}(\overline{Q}, \delta) \rightarrow \mathfrak{g}^*$ defined in the introduction. We will denote by $\mu^{-1}(0)$ the scheme-theoretic fiber of μ over 0. Thus

$$\mathbb{C}[\mu^{-1}(0)] = \frac{\mathbb{C}[\text{Rep}(\overline{Q}, \delta)]}{\mathbb{C}[\text{Rep}(\overline{Q}, \delta)] \cdot \mathfrak{g}}, \quad (2.1)$$

where, by abuse of notation, $\mathfrak{g}$ denotes the linear subspace of $\mathbb{C}[\text{Rep}(\overline{Q}, \delta)]$ obtained by pulling back via μ the elements of $\mathfrak{g}$ viewed as linear functions on $\mathfrak{g}^*$. However, by a result of Crawley-Boevey [C-B], the scheme $\mu^{-1}(0)$ is in fact reduced and irreducible. (The reason for defining $\mu^{-1}(0)$ as the scheme-theoretic fiber is that (2.1) will be important for us later on.) We define

$$R = \mathbb{C}[\mu^{-1}(0)]^G.$$

It is well known that

$$\text{Spec } R = \text{Spec } \mathbb{C}[\mu^{-1}(0)]^G = \mu^{-1}(0)//G \cong X. \quad (2.2)$$

In particular, R is a normal, 2-dimensional, commutative Gorenstein domain.

2.2. Cassens and Slodowy ([CS], §7) explain that a minimal resolution of X can be constructed as a GIT quotient

$$\tilde{X} = \mu^{-1}(0)_{\chi}^{ss}/G \quad (2.3)$$

for any $\chi \in \Lambda_{++}$, where $\mu^{-1}(0)_{\chi}^{ss}$ denotes the open subset of $\mu^{-1}(0)$ consisting of the points *semistable* with respect to χ . Moreover, they prove that for each such χ ,

- (A) $\mu^{-1}(0)_{\chi}^{ss} = \mu^{-1}(0)_{\chi}^s$, the set of *stable* points with respect to χ , and
- (B) the action of G on $\mu^{-1}(0)_{\chi}^s$ is *free* (recall that, a priori, the action of G on the set of stable points only needs to have finite stabilizers; in our situation, however, all stabilizers turn out to be trivial).

Moreover, (2.2) and (2.3) lead to the description of the resolution $\tilde{X} \rightarrow X$ as the natural map

$$\mathrm{Proj} S \longrightarrow \mathrm{Spec} S_0,$$

where S is the graded ring

$$S = \bigoplus_{n \geq 0} T^{G, \chi^n},$$

where $T = \mathbb{C}[\mu^{-1}(0)]$.

Lemma 2.1. *The ring S is finitely generated.*

Proof. This is a very general statement. Write $Y = \mathrm{Spec} T$, and consider the induced action of G on $Y \times \mathbb{C}$, where the action on the first factor is the given one, and the action on $\mathbb{C}$ is via χ . The ring $\mathbb{C}[Y \times \mathbb{C}] = T \otimes \mathbb{C}[z]$ has the obvious grading by the degree of polynomials with respect to z , and we clearly have an isomorphism of graded algebras

$$\mathbb{C}[Y \times \mathbb{C}]^G \cong \bigoplus_{n \geq 0} T^{G, \chi^n}.$$

In particular, the algebra on the RHS is finitely generated (here we have used the fact that G is reductive). $\square$

2.3. The main goal of this section is to obtain some more detailed information on the ring S , in the form of the following result.

Theorem 2.2. *Let $p : \mu^{-1}(0)_\chi^{ss} \rightarrow \tilde{X}$ denote the quotient map.*

- (1) *There exists a unique line bundle $\mathcal{L}$ on $\tilde{X}$ such that $p^*\mathcal{L}$ is the trivial line bundle on $\mu^{-1}(0)_\chi^{ss}$ equipped with the G -linearization given by the character χ . Moreover, $\mathcal{L}$ is ample.*
- (2) *The induced map*

$$S_n = \mathbb{C}[\mu^{-1}(0)]^{G, \chi^n} \longrightarrow \Gamma(\tilde{X}, \mathcal{L}^{\otimes n})$$

is an isomorphism for sufficiently large n . In particular, S_n is a torsion-free S_0 -module of generic rank 1 for sufficiently large n .

- (3) *The multiplication map*

$$S_m \otimes S_n \longrightarrow S_{m+n}$$

is surjective for sufficiently large m and n .

2.4. It will be clear from the proof of the theorem that essentially the only properties that we use are the fact that G is reductive, statements (A) and (B) of §2.2, and the fact that $\mu^{-1}(0)_\chi^{ss}$ is dense in $\mu^{-1}(0)$. Thus the theorem could be stated and proved in a much more general context, where $\mu^{-1}(0)$ is replaced by any affine variety Y with an action of G satisfying properties (A) and (B) of §2.2, such that the set of semistable points Y_χ^{ss} is dense in Y .

2.5. We begin the proof of Theorem 2.2 by observing that (A) and (B) imply that the quotient map p is a principal G -bundle. Now it is easy to see that the notion of a G -linearization for a coherent sheaf on $\mu^{-1}(0)_\chi^{ss}$ is equivalent to the notion of a descent datum for the (flat) morphism p . Hence the first statement of part (1) of the theorem follows immediately from flat descent theory. Another consequence of descent theory is that for any line bundle $\mathcal{M}$ on $\tilde{X}$, we have

$$\Gamma(\tilde{X}, \mathcal{M}) = \Gamma(\mu^{-1}(0)_\chi^{ss}, p^* \mathcal{M})^G. \quad (2.4)$$

Indeed, the LHS of (2.4) coincides with $\text{Hom}(\mathcal{O}_{\tilde{X}}, \mathcal{M})$. But $p^* \mathcal{O}_{\tilde{X}}$ is the trivial line bundle on $\mu^{-1}(0)_\chi^{ss}$ equipped with the trivial G -linearization, and descent theory for morphisms implies that

$$\text{Hom}(\mathcal{O}_{\tilde{X}}, \mathcal{M}) = \text{Hom}_{G\text{-equiv}}(p^* \mathcal{O}_{\tilde{X}}, p^* \mathcal{M}),$$

which proves (2.4).

2.6. The construction of the quotient (2.3) given in [GIT] shows that for some $N \in \mathbb{N}$, there exists an ample line bundle $\mathcal{L}'$ on $\tilde{X}$ such that $p^* \mathcal{L}'$ is the trivial line bundle on $\mu^{-1}(0)_\chi^{ss}$ equipped with the G -linearization given by χ^N . Moreover, $\mathcal{L}' = \mathcal{O}_{\tilde{X}}(N)$ for the description of $\tilde{X}$ as $\text{Proj } S$. Now the discussion in §2.5 implies that $\mathcal{L}' = \mathcal{L}^{\otimes N}$; in particular, $\mathcal{L}$ itself is ample, completing the proof of part (1) of the theorem.

Replacing N by one of its multiples if necessary, we may assume that

$$S_{jN} = (S_N)^j \quad \text{for all } j \geq 1;$$

this follows from the fact that S is finitely generated (Lemma 2.1) and Lemma 2.1.6(v) of [EGA II]. Similarly, we may assume that $\mathcal{L}^{\otimes N}$ is very ample (using Proposition 4.5.10(ii) of loc. cit.). And, finally, we may assume that the natural map

$$S_{jN} \longrightarrow \Gamma(\tilde{X}, \mathcal{L}^{\otimes jN})$$

is an isomorphism for all $j \geq 1$. From now on we fix $N \in \mathbb{N}$ satisfying all the properties listed above.

2.7. The line bundle $\mathcal{L}^{\otimes n}$ is very ample for all sufficiently large n (loc. cit., Proposition 4.5.10(ii)). In particular, there exists $d_1 \in \mathbb{N}$ such that each of the bundles

$$\mathcal{L}^{\otimes d_1 N}, \mathcal{L}^{\otimes (d_1 N + 1)}, \dots, \mathcal{L}^{\otimes (d_1 N + N - 1)} \quad (2.5)$$

is generated by global sections. But for any $n \in \mathbb{N}$, we have, from (2.4),

$$\Gamma(\tilde{X}, \mathcal{L}^{\otimes n}) = \Gamma(\mu^{-1}(0)_\chi^{ss}, \mathcal{O})^{G, \chi^n}.$$

Now recall that $\mu^{-1}(0)_\chi^{ss}$ is precisely the set of points of $\mu^{-1}(0)$ where at least one element of $\mathbb{C}[\mu^{-1}(0)]^{G, \chi^N} = S_N$ does not vanish. In particular, if $\sigma \in \Gamma(\tilde{X}, \mathcal{L}^{\otimes n})$, then there exist $j \in \mathbb{N}$ and finitely many elements $f_1, \dots, f_r \in S_{jN}$ such that $f_i \sigma \in S_{n+jN}$ for each i , and the open sets $\{f_i \neq 0\}$ cover all of $\mu^{-1}(0)_\chi^{ss}$.

Since we are dealing with finitely many line bundles (2.5), we deduce that there exists $d_2 \in \mathbb{N}$ such that for every $0 \leq k \leq N - 1$, the line bundle $\mathcal{L}^{\otimes(d_1N+d_2N+k)}$ is generated by finitely many sections coming from the elements of $S_{d_1N+d_2N+k}$.

For every $0 \leq k \leq N - 1$, let us now choose a finite dimensional subspace

$$V_k \subseteq S_{d_1N+d_2N+k} \subseteq \Gamma(\tilde{X}, \mathcal{L}^{\otimes(d_1N+d_2N+k)})$$

of sections which generate the line bundle $\mathcal{L}^{\otimes(d_1N+d_2N+k)}$. These sections determine a surjection of coherent sheaves

$$\phi_k : \mathcal{O}_{\tilde{X}} \otimes_{\mathbb{C}} V_k \longrightarrow \mathcal{L}^{\otimes(d_1N+d_2N+k)}. \quad (2.6)$$

Let $\mathcal{N}_k$ denote the kernel of this surjection; it is a coherent sheaf on $\tilde{X}$. Since $\mathcal{L}^{\otimes N}$ is very ample, there exists $d_3 \in \mathbb{N}$ such that

$$H^1(\tilde{X}, \mathcal{L}^{\otimes jN} \otimes_{\mathcal{O}_{\tilde{X}}} \mathcal{N}_k) = 0 \quad (2.7)$$

for every $j \geq d_3$ and every $0 \leq k \leq N - 1$.

2.8. We can now prove part (2) of Theorem 2.2. Namely, every integer $n \geq (d_1 + d_2 + d_3) \cdot N$ can be written as $n = jN + d_1N + d_2N + k$ for some (uniquely determined) $j \geq d_3$ and $0 \leq k \leq N - 1$. We have a short exact sequence, induced by (2.6):

$$0 \longrightarrow \mathcal{L}^{\otimes jN} \otimes_{\mathcal{O}_{\tilde{X}}} \mathcal{N}_k \longrightarrow \mathcal{L}^{\otimes jN} \otimes_{\mathbb{C}} V_k \longrightarrow \mathcal{L}^{\otimes n} \longrightarrow 0.$$

Applying the long exact cohomology sequence and using (2.7), we see that the map

$$\Gamma(\tilde{X}, \mathcal{L}^{\otimes jN} \otimes_{\mathbb{C}} V_k) \longrightarrow \Gamma(\tilde{X}, \mathcal{L}^{\otimes n})$$

is surjective. But

$$\Gamma(\tilde{X}, \mathcal{L}^{\otimes jN} \otimes_{\mathbb{C}} V_k) = \Gamma(\tilde{X}, \mathcal{L}^{\otimes jN}) \otimes_{\mathbb{C}} V_k = S_{jN} \otimes_{\mathbb{C}} V_k \subseteq S_{jN} \otimes_{\mathbb{C}} S_{d_1N+d_2N+k},$$

and so, a fortiori, the natural map

$$S_n \longrightarrow \Gamma(\tilde{X}, \mathcal{L}^{\otimes n})$$

is surjective for all $n \geq (d_1 + d_2 + d_3) \cdot N$. Furthermore, this map is injective for all n because $\mu^{-1}(0)_{\chi}^{ss}$ is dense in $\mu^{-1}(0)$ (since $\mu^{-1}(0)$ is irreducible). Observe moreover that $\Gamma(\tilde{X}, \mathcal{L}^{\otimes n})$ is a finitely generated $\mathbb{C}[\mu^{-1}(0)]$ -module of generic rank 1, since $\tilde{X} \rightarrow X$ is a projective birational map which is an isomorphism away from the fiber over the singular point $0 \in X$. This proves part (2) of the theorem.

2.9. Finally, the proof of part (3) of Theorem 2.2 is a variation of the idea used above, with an extra trick. It is best to formulate it as a general statement. In view of parts (1) and (2), it is clear that part (3) follows from the following

Proposition 2.3. *Let Y be a scheme, projective over a (commutative) Noetherian ring A , and let $\mathcal{L}$ be an ample invertible sheaf on Y . Then the natural map*

$$\Gamma(Y, \mathcal{L}^{\otimes m}) \otimes_A \Gamma(Y, \mathcal{L}^{\otimes n}) \longrightarrow \Gamma(Y, \mathcal{L}^{\otimes(m+n)}) \quad (2.8)$$

is surjective for all sufficiently large m and n .

Proof. As before, the sheaves $\mathcal{L}^{\otimes n}$ are very ample for all $n \gg 0$. In particular, there exists $N \in \mathbb{N}$ such that for every $N \leq k \leq 2N-1$, the sheaf $\mathcal{L}^{\otimes k}$ is generated by a finitely generated A -submodule of global sections $V_k \subseteq \Gamma(Y, \mathcal{L}^{\otimes k})$. Let us write $\phi_k : \mathcal{O}_Y \otimes_A V_k \rightarrow \mathcal{L}^{\otimes k}$ for the corresponding surjection, and $\mathcal{N}_k$ for its kernel.

Next, let $N' \in \mathbb{N}$ be such that

$$H^1(Y, \mathcal{L}^{\otimes n} \otimes_{\mathcal{O}_Y} \mathcal{N}_k) = 0 \quad \text{for all } n \geq N' \text{ and } N \leq k \leq 2N-1.$$

We claim that if $n \geq N'$ and $m \geq 2N-1$, then the map (2.8) is surjective. Indeed, if $m \geq 2N-1$ is an integer, it can clearly be written as a sum of the form $m = k_1 + \cdots + k_r$, where $r \geq 1$ and $N \leq k_j \leq 2N-1$ for every j . Now for $0 \leq j \leq r-1$, we have an exact sequence

$$0 \longrightarrow \mathcal{L}^{\otimes(n+k_1+\cdots+k_j)} \otimes_{\mathcal{O}_Y} \mathcal{N}_k \longrightarrow \mathcal{L}^{\otimes(n+k_1+\cdots+k_j)} \otimes_A V_{k_{j+1}} \longrightarrow \mathcal{L}^{\otimes(n+k_1+\cdots+k_j+k_{j+1})} \longrightarrow 0,$$

and since $n + k_1 + \cdots + k_j \geq n \geq N'$, we see that the induced map

$$\Gamma(Y, \mathcal{L}^{\otimes(n+k_1+\cdots+k_j)}) \otimes_A V_{k_{j+1}} \longrightarrow \Gamma(Y, \mathcal{L}^{\otimes(n+k_1+\cdots+k_j+k_{j+1})})$$

is surjective. A fortiori, this implies that the map

$$\Gamma(Y, \mathcal{L}^{\otimes(n+k_1+\cdots+k_j)}) \otimes_A \Gamma(Y, \mathcal{L}^{\otimes k_{j+1}}) \longrightarrow \Gamma(Y, \mathcal{L}^{\otimes(n+k_1+\cdots+k_j+k_{j+1})})$$

is surjective.

By induction, we deduce that the map

$$\Gamma(Y, \mathcal{L}^{\otimes n}) \otimes_A \Gamma(Y, \mathcal{L}^{\otimes k_1}) \otimes_A \cdots \otimes_A \Gamma(Y, \mathcal{L}^{\otimes k_r}) \longrightarrow \Gamma(Y, \mathcal{L}^{\otimes(m+n)})$$

is surjective. A fortiori, this implies that the map (2.8) is surjective, completing the proof of the proposition (and thereby of Theorem 2.2). $\square$

3. QUANTIZATION OF KLEINIAN SINGULARITIES

3.1. In this section we recall some results of M.P. Holland [Ho] that are crucial for our construction of the quantization of the minimal resolution. We will use the notation

$$\Gamma, I, Q, \delta, G, \mathfrak{g}, \mu, \text{ etc.}$$

defined in the introduction. If $a \in Q$ is an arrow, we write $t(a), h(a) \in I$ for the tail and head of a , respectively. The *defect* $\partial \in \mathbb{Z}^I$ is defined by

$$\partial_i = -\delta_i + \sum_{t(a)=i} \delta_{h(a)} \quad \text{for all } i \in I.$$

We identify $\mathbb{C}_0^I := \{\chi \in \mathbb{C}^I \mid \chi \cdot \delta = 0\}$ with the space of 1-dimensional characters of $\mathfrak{g}$ via the various trace maps $\mathfrak{gl}(\delta_i, \mathbb{C}) \rightarrow \mathbb{C}$. If $\chi \in \mathbb{C}_0^I$, we define a filtered algebra

$$\mathcal{U}^\chi = \frac{\mathcal{D}(\text{Rep}(Q, \delta))^G}{\left[\mathcal{D}(\text{Rep}(Q, \delta)) \cdot (\iota - \chi)(\mathfrak{g}) \right]^G},$$

where $\mathcal{D}(\text{Rep}(Q, \delta))$ is the algebra of polynomial differential operators on the affine space $\text{Rep}(Q, \delta)$, and

$$\iota : \mathfrak{g} \longrightarrow \text{Vect}(\text{Rep}(Q, \delta)) \subset \mathcal{D}(\text{Rep}(Q, \delta))$$

is the Lie algebra map induced by the G -action.

CAUTION. For consistency with the filtration on the algebras $\mathcal{O}^\lambda$ introduced in §1.1, we need to use the Bernstein filtration on the algebra $\mathcal{D}(\text{Rep}(Q, \delta))$ (instead of the more standard order filtration). Fortunately, as remarked in [Ho], the results of sections 2–4 of that paper remain valid if the order filtration is replaced by the Bernstein filtration. From now on it will be implicitly assumed that the results of all constructions involving differential operators will be equipped with filtrations induced from the Bernstein filtration.

Theorem 3.1 (Holland). *If $\lambda \in \mathbb{C}^I$ is such that $\lambda \cdot \delta = 1$, then there is a natural isomorphism of filtered algebras*

$$\mathcal{O}^\lambda \cong \mathcal{U}^{\lambda - \partial - \epsilon_0}$$

where $\epsilon_0 \in \mathbb{Z}^I$ is the standard basis vector corresponding to the extending vertex.

Proof. See [Ho], Corollary 4.7. □

3.2. The following result will also be important to us.

Theorem 3.2. *If $\chi \in \mathbb{C}_0^I$, there is a natural isomorphism*

$$\text{gr}^\bullet \left(\frac{\mathcal{D}(\text{Rep}(Q, \delta))}{\mathcal{D}(\text{Rep}(Q, \delta)) \cdot (\iota - \chi)(\mathfrak{g})} \right) \cong \frac{\text{gr}^\bullet \mathcal{D}(\text{Rep}(Q, \delta))}{[\text{gr}^\bullet \mathcal{D}(\text{Rep}(Q, \delta))] \cdot \mathfrak{g}}$$

of graded $\mathbb{C}[T^* \text{Rep}(Q, \delta)]$ -modules.

Proof. Combine [Ho], Proposition 2.4 with the fact that the moment map $\mu : \text{Rep}(\overline{Q}, \delta) \rightarrow \mathfrak{g}^*$ is flat ([CBH], Lemma 8.3). □

4. MORITA $\mathbb{Z}$ -ALGEBRAS

4.1. In this section we review the basic theory of $\mathbb{Z}$ -algebras following [GS]. We also give a detailed proof of a strengthening of Lemma 5.5 of loc. cit. that is used in our paper.

Definition 4.1. A lower-triangular $\mathbb{Z}$ -algebra is given by the following collection of data.

(a) An abelian group B , bigraded by $\mathbb{Z}$ in the following way:

$$B = \bigoplus_{i \geq j \geq 0} B_{ij}.$$

(b) An algebra structure on each of the additive groups $B_i := B_{ii}$.

(c) A (B_i, B_j) -bimodule structure on B_{ij} for each $i > j \geq 0$.

(d) A map of (B_i, B_k) -bimodules

$$B_{ij} \otimes_{B_j} B_{jk} \longrightarrow B_{ik}$$

for each $i > j > k \geq 0$.

The multiplication on the additive group B is defined so as to imitate multiplication of lower-triangular matrices in the obvious sense, and the maps in (d) are required to satisfy the obvious compatibility conditions needed to make the multiplication on B associative. Recall that the algebras B_i are assumed to have multiplicative identities; however, B will almost never have a multiplicative identity, since it is defined as an infinite direct sum.

Next we consider modules over $\mathbb{Z}$ -algebras.

Definition 4.2. Let B be a lower-triangular $\mathbb{Z}$ -algebra as in the definition above. A *graded B -module* is given by the following collection of data.

(a) A positively graded abelian group

$$M = \bigoplus_{i \geq 0} M_i.$$

(b) A B_i -module structure on M_i for each $i \geq 0$.

(c) A map of left B_i -modules

$$B_{ij} \otimes_{B_j} M_j \longrightarrow M_i$$

for each $i > j \geq 0$.

The action of B on M is defined so as to imitate multiplication of column vectors by lower-triangular matrices on the left, and the maps in (c) are required to satisfy the obvious compatibility conditions needed to make M into a left B -module. Note also that for each $i \geq 0$, the element $1 \in B_i$ is assumed to act as the identity on M_i .

4.2. With these definitions in hand, we construct the categories $B\text{-Grmod}$, $B\text{-Tors}$, $B\text{-Qgr}$, $B\text{-grmod}$, $B\text{-tors}$ and $B\text{-qgr}$ in the same way as for graded rings. If $A = \bigoplus_{n \geq 0} A_n$ is a graded ring, we can associate to it a lower-triangular $\mathbb{Z}$ -algebra $B = \hat{A}$ by defining $B_{ij} = A_{i-j}$ for $i \geq j \geq 0$. As explained in §5.3 of [GS], we then have natural equivalences of categories

$$A\text{-Qgr} \xrightarrow{\sim} \hat{A}\text{-Qgr} \quad \text{and} \quad A\text{-qgr} \xrightarrow{\sim} \hat{A}\text{-qgr}.$$

4.3. We are now ready for the key definition; note that it is *weaker* than the corresponding notion introduced in §5.4 of [GS].

Definition 4.3. A *Morita $\mathbb{Z}$ -algebra* is a lower-triangular $\mathbb{Z}$ -algebra

$$B = \bigoplus_{i \geq j \geq 0} B_{ij}$$

such that there exists $N \in \mathbb{N}$ for which

- (i) the (B_i, B_j) -bimodule B_{ij} yields an equivalence

$$B_j\text{-mod} \xrightarrow{\sim} B_i\text{-mod}$$

whenever $i - j \geq N$; and

- (ii) the multiplication map

$$B_{ij} \otimes_{B_j} B_{jk} \longrightarrow B_{ik}$$

is an isomorphism whenever $i - j, j - k \geq N$.

Theorem 4.4. *Suppose that B is a Morita $\mathbb{Z}$ -algebra such that each B_i is a left Noetherian ring, and each B_{ij} is a finitely generated left B_i -module. Then*

- (1) *Each finitely generated graded left B -module is graded-Noetherian.*
- (2) *The association*

$$\phi : M \longmapsto \bigoplus_{n \geq 0} B_{n,0} \otimes_{B_0} M$$

induces an equivalence of categories

$$\Phi : B_0\text{-mod} \xrightarrow{\sim} B\text{-qgr}.$$

Proof. We follow the proof of Lemma 5.5 in [GS] closely.

(1) Let M be a finitely generated graded B -module. We have to show that every graded submodule of M is also finitely generated. It is clear that M is generated by finitely many homogeneous elements, so it is enough to consider the case where M is generated by one homogeneous element, say of degree a . In this case M is a graded homomorphic image of $\bigoplus_{j \geq a} B_{ja}$, so we assume, without loss of generality, that

$$M = \bigoplus_{j \geq a} B_{ja}.$$

Now let

$$L = \bigoplus_{j \geq a} L_j \subseteq M$$

be a graded submodule. We use the notation

$$B_{ij}^* = \text{Hom}_{B_i\text{-mod}}(B_{ij}, B_i),$$

which is a (B_j, B_i) -bimodule. Let $N \in \mathbb{N}$ be as in the definition of a Morita $\mathbb{Z}$ -algebra. Then for $j \geq a + N$, we have a chain of maps of left B_a -modules

$$B_{ja}^* \otimes_{B_j} L_j \hookrightarrow B_{ja}^* \otimes_{B_j} M_j = B_{ja}^* \otimes_{B_j} B_{ja} \xrightarrow{\cong} B_a,$$

where the first map is injective because B_{ja}^* is a projective right B_j -module, and the second map is an isomorphism by definition. We let

$$X(j) \subseteq B_a$$

be the image of the composition above. Since B_a is assumed to be Noetherian, there exists an integer $b \geq a + N$ such that

$$\sum_{j \geq a+N} X(j) = \sum_{i=a+N}^b X(i) \subseteq B_a.$$

Now for $k \geq a + N$, we have

$$L_k \cong B_{ka} \otimes_{B_a} B_{ka}^* \otimes_{B_k} L_k,$$

which means that

$$L_k = B_{ka} X(k) \quad \text{for } k \geq a + N,$$

as submodules of $B_{ka} = M_k$. Thus, for $k \geq b + N$, we have

$$\begin{aligned} L_k &= B_{ka} X(k) \subseteq \sum_{i=a+N}^b B_{ka} X(i) \\ &= \sum_{i=a+N}^b B_{ki} B_{ia} X(i) = \sum_{i=a+N}^b B_{ki} L_i, \end{aligned}$$

where we have used the assumption that $B_{ki} \otimes_{B_i} B_{ia} \xrightarrow{\cong} B_{ka}$ for $k - i, i - a \geq N$. Thus we see that

$$\sum_{k \geq b+N} L_k \text{ is generated by } \sum_{i=a+N}^b L_i$$

as a B -module. Finally, for $a \leq j \leq b + N$, L_j is a B_j -submodule of the finitely generated B_j -module $M_j = B_{ja}$, and is therefore finitely generated, completing the proof of (1).

(2) Note that, in view of (1), the category B -tors consists precisely of the bounded finitely generated graded left B -modules. This remark will be used without further mention throughout the proof.

We construct a functor

$$\Theta : B\text{-qgr} \longrightarrow B_0\text{-mod}$$

as follows. Let $\widetilde{M} \in B\text{-qgr}$ be the image of some $M \in B\text{-grmod}$. Then M is generated by $\bigoplus_{i=0}^a M_i$ as a left B -module, for some $a \in \mathbb{N}$. Thus

$$M_j = \sum_{i=0}^a B_{ji} M_i \quad \text{for each } j \geq a,$$

and so, by assumption,

$$M_k = \sum_{i=0}^a B_{ki} M_i = \sum_{i=0}^a B_{kj} B_{ji} M_i = B_{kj} M_j$$

whenever $j \geq a$ and $k \geq j + N$. So if $k - j, j - a \geq N$, we have a chain of natural homomorphisms

$$\begin{aligned} B_{ja}^* \otimes_{B_j} M_j &\xrightarrow{\simeq} B_{ja}^* \otimes_{B_j} B_{kj}^* \otimes_{B_k} B_{kj} \otimes_{B_j} M_j \\ &\xrightarrow{\simeq} B_{ka}^* \otimes_{B_k} (B_{kj} \otimes_{B_j} M_j) \twoheadrightarrow B_{ka}^* \otimes_{B_k} M_k. \end{aligned}$$

Let θ_{kj} denote the composition, which is a surjective homomorphism of left B_a -modules:

$$\theta_{kj} : B_{ja}^* \otimes_{B_j} M_j \twoheadrightarrow B_{ka}^* \otimes_{B_k} M_k.$$

Since the construction is natural, it is clear that the homomorphisms θ_{kj} satisfy the compatibility conditions

$$\theta_{lj} = \theta_{lk} \circ \theta_{kj} \quad \text{for all } l - k, k - j, j - a \geq N.$$

Sublemma 4.5. *There exists an integer $b \geq a + N$ such that the map θ_{kj} is an isomorphism whenever $j \geq b$ and $k - j \geq N$.*

Suppose otherwise. Then there exist $j_1 \geq a + N$ and $k_1 \geq j_1 + N$ such that $\theta_{k_1 j_1}$ has nonzero kernel. Further, there exist $j_2 \geq k_1 + N$ and $k_2 \geq j_2 + N$ such that $\theta_{k_2 j_2}$ has nonzero kernel. Continuing this process inductively, we see that there exists an infinite sequence of surjective homomorphisms

$$B_{j_1 a}^* \otimes_{B_{j_1}} M_{j_1} \xrightarrow{\theta_{k_1 j_1}} B_{k_1 a}^* \otimes_{B_{k_1}} M_{k_1} \xrightarrow{\theta_{j_2 k_1}} B_{j_2 a}^* \otimes_{B_{j_2}} M_{j_2} \xrightarrow{\theta_{k_2 j_2}} \dots$$

that never stabilizes to a sequence of isomorphisms. But since $B_{j_1 a}^*$ is a finitely generated left B_a -module and M_{j_1} is a finitely generated left B_{j_1} -module, we see that $B_{j_1 a}^* \otimes_{B_{j_1}} M_{j_1}$ is a finitely generated, hence Noetherian, left B_a -module, which is a contradiction.

We now fix $b \in \mathbb{N}$ satisfying the conditions of the sublemma. We define

$$L = B_{b0}^* \otimes_{B_b} M_b \in B_0\text{-mod}.$$

Sublemma 4.6. *We have $\Phi(L) \cong \widetilde{M}$ in $B\text{-qgr}$, where the functor Φ is defined in the statement of the theorem.*

Recall that $\Phi(L)$ is the image under the quotient map of

$$\phi(L) = \bigoplus_{n \geq 0} B_{n0} \otimes_{B_0} B_{b0}^* \otimes_{B_b} M_b.$$

Note that for $n \geq b + N$, we have the natural isomorphism of (B_n, B_0) -bimodules

$$B_{nb} \otimes_{B_b} B_{b0} \xrightarrow{\simeq} B_{n0}.$$

Tensoring with B_{b0}^* on the right, we obtain a natural isomorphism

$$B_{nb} \xrightarrow{\simeq} B_{n0} \otimes_{B_0} B_{b0}^*.$$

Similarly, we have

$$B_{kj} \xrightarrow{\simeq} B_{ka} \otimes_{B_a} B_{ja}^*$$

for $j \geq b$, $k \geq j + N$. Thus we obtain natural isomorphisms

$$M_k \xrightarrow{\simeq} B_{ka} \otimes_{B_a} B_{ka}^* \otimes_{B_k} M_k \xrightarrow{(1 \otimes \theta_{kj})^{-1}} B_{ka} \otimes_{B_a} B_{ja}^* \otimes_{B_j} M_j \xrightarrow{\simeq} B_{kj} \otimes_{B_j} M_j.$$

In particular, taking $j = b$ and combining with the previous remarks, we obtain natural isomorphisms

$$B_{n0} \otimes_{B_0} B_{b0}^* \otimes_{B_b} M_b \xrightarrow{\simeq} B_{nb} \otimes_{B_b} M_b \xrightarrow{\simeq} M_n$$

for all $n \geq b + N$. By the naturality of all the constructions, we see that we have defined an isomorphism of graded left B -modules

$$\phi(L)_{\geq b+N} \xrightarrow{\simeq} M_{\geq b+N} = \bigoplus_{n \geq b+N} M_n,$$

completing the proof of the sublemma.

We note that, by similar arguments, replacing b with any $b' \geq b$ yields a module $L' = B_{b'0}^* \otimes_{B_{b'}} M_{b'} \in B_0\text{-mod}$ that is canonically isomorphic to L . In particular, L only depends on the image $\widetilde{M}$ of M in $B\text{-qgr}$, up to isomorphism.

Going back to the construction of the functor

$$\Theta : B\text{-qgr} \longrightarrow B_0\text{-mod},$$

we set $\Theta(\widetilde{M}) = L$ with the notation above. We need to explain how Θ acts on morphisms. Suppose $M, M' \in B\text{-grmod}$ and $L = \Theta(\widetilde{M})$, $L' = \Theta(\widetilde{M}')$. In the construction of L , L' given above, we pick $b \in \mathbb{N}$ that works for both; thus $L = B_{b0}^* \otimes_{B_b} M_b$ and $L' = B_{b0}^* \otimes_{B_b} M'_b$. Now a morphism of graded B -modules

$$f : M \longrightarrow M'$$

gives in particular a B_b -module homomorphism $f_b : M_b \rightarrow M'_b$; tensoring with B_{b0}^* yields a B_0 -module morphism

$$\Theta(f) := 1 \otimes f_b : L \longrightarrow L'.$$

Finally, we need to check that the functors Φ and Θ are quasi-inverse to each other. It is essentially obvious that $\Theta \circ \Phi$ is isomorphic to the identity functor on $B_0\text{-mod}$, so we only need to consider the composition $\Phi \circ \Theta$.

In the proof of Sublemma 4.6 we have constructed a natural isomorphism

$$\Phi(\Theta(\widetilde{M})) \xrightarrow{\sim} \widetilde{M}$$

for every $M \in B\text{-grmod}$. We only need to see that this isomorphism is compatible with morphisms; the essential observation is that if $f : M \rightarrow M'$ is a morphism in $B\text{-grmod}$ and b is as in the previous paragraph, then f_n is uniquely determined by f_b for all $n \geq b + N$, due to the commutativity of the diagram

$$\begin{array}{ccc} B_{nb} \otimes_{B_b} M_b & \xrightarrow{1 \otimes f_b} & B_{nb} \otimes_{B_b} M'_b \\ \downarrow & & \downarrow \\ M_n & \xrightarrow{f_n} & M'_n \end{array}$$

where the vertical arrows are isomorphisms. This completes the proof of the theorem. $\square$

5. QUANTIZATION OF THE MINIMAL RESOLUTION

5.1. In this section we use Holland's results described in Section 3 to define the algebras $B^\lambda(\chi)$ mentioned in the introduction and state our main result. We use the same notation as in the introduction and in Section 3. We let $\mathfrak{X}(G)$ denote the group of the algebraic group homomorphisms $\zeta : G \rightarrow \mathbb{C}^\times$. Note that if $\zeta \in \mathfrak{X}(G)$, its differential $d\zeta : \mathfrak{g} \rightarrow \mathbb{C}$ can be thought of as an element of $\mathbb{C}_0^I$ (see Section 3).

Given $\zeta \in \mathfrak{X}(G)$ and $\chi \in \mathbb{C}_0^I$, we define

$$P_{\chi, \zeta} = \frac{\mathcal{D}(\text{Rep}(Q, \delta))^{G, \zeta}}{\left[\mathcal{D}(\text{Rep}(Q, \delta)) \cdot (\iota - \chi)(\mathfrak{g}) \right]^{G, \zeta}}.$$

One immediately checks that the actions of $\mathcal{D}(\text{Rep}(Q, \delta))^G$ on $\mathcal{D}(\text{Rep}(Q, \delta))^{G, \zeta}$ by left and right multiplication descend to a filtered $(\mathcal{U}^{\chi+d\zeta}, \mathcal{U}^\chi)$ -bimodule structure on $P_{\chi, \zeta}$.

5.2. If $\lambda \in \mathbb{C}^I$ is such that $\lambda \cdot \delta = 1$, we will write

$$P_\zeta^\lambda = P_{\lambda - \partial - \epsilon_0, \zeta}.$$

According to Theorem 3.1, we can think of P_ζ^λ as a filtered $(\mathcal{O}^{\lambda+d\zeta}, \mathcal{O}^\lambda)$ -bimodule. Moreover, by Theorem 3.2, we have an isomorphism of graded bimodules

$$\text{gr}^\bullet P_\zeta^\lambda \cong \mathbb{C}[\mu^{-1}(0)]^{G, \zeta}. \quad (5.1)$$

The following fact will be used implicitly in Section 6; it is needed in order to justify the use of Proposition 6.9(1).

Lemma 5.1. *The filtration on P_ζ^λ induced by the Bernstein filtration on differential operators is good in the sense of [Bj], Definition 2.19.*

Proof. In view of (5.1) and the Remark after the proof of Proposition 2.22 in [Bj], it suffices to show that $\mathbb{C}[\mu^{-1}(0)]^{G, \zeta}$ is a finitely generated $\mathbb{C}[\mu^{-1}(0)]^G$ -module for every ζ . Now the algebra $\oplus_{n \geq 0} \mathbb{C}[\mu^{-1}(0)]^{G, \zeta^n}$ is finitely generated by the argument given in the proof of Lemma 2.1, and therefore our claim follows from Lemma 2.1.6(i) in [EGA II]. $\square$

5.3. Observe now that differentiation of characters induces an isomorphism of abelian groups $d : \mathfrak{X}(G) \xrightarrow{\sim} \Lambda \subseteq \mathbb{C}_0^I$; by abuse of notation, if $\xi \in \Lambda$, we will write

$$P_\xi^\lambda = P_\zeta^\lambda \quad \text{and} \quad \mathbb{C}[\mu^{-1}(0)]^{G, \xi} = \mathbb{C}[\mu^{-1}(0)]^{G, \zeta}$$

where $\zeta \in \mathfrak{X}(G)$ is such that $d\zeta = \xi$. Now, given $\chi \in \Lambda_{++}$, we define a lower-triangular $\mathbb{Z}$ -algebra $B(\lambda, \chi)$ by

$$B(\lambda, \chi)_{ij} = \begin{cases} \mathcal{O}^{\lambda+j\cdot\chi} & \text{if } i = j \geq 0, \\ P_{(i-j)\cdot\chi}^{\lambda+j\cdot\chi} & \text{if } i > j \geq 0. \end{cases}$$

All the structure maps of this $\mathbb{Z}$ -algebra are induced by the multiplication of elements of $\mathcal{D}(\text{Rep}(Q, \delta))^G$, and all compatibility conditions follow immediately from the associativity of this multiplication. Observe that $B(\lambda, \chi)$ is naturally filtered by the Bernstein filtration of differential operators. (We leave the formulation of the general notion of a filtered $\mathbb{Z}$ -algebra B to the reader: each component B_{ij} should be positively filtered, and all the structure maps should be compatible with the filtrations. See also [GS2].)

5.4. We are now ready to state our main result, whose proof occupies Section 6.

Theorem 5.2. *Let $\lambda \in \mathbb{C}^I$ be such that $\lambda \cdot \delta = 1$ and $\lambda \cdot \alpha \neq 0$ for every Dynkin root α (i.e., the algebra $\mathcal{O}^\lambda$ is regular). Given $\chi \in \Lambda_{++}$, there exists $\xi \in \Lambda_{++}$ such that the lower-triangular filtered $\mathbb{Z}$ -algebra*

$$B^\lambda(\chi) := B(\lambda + \xi, \chi),$$

where $B(\lambda + \xi, \chi)$ is constructed above, satisfies the properties (1) and (2) of §1.5 of the introduction.

This theorem answers a question posed by M.P. Holland on page 831 of [Ho]. He constructs a quantization of a certain *partial* resolution of the Kleinian singularity $X = \mathbb{C}^2/\Gamma$, and asks for a quantization of the whole minimal resolution. It appears that the formalism of $\mathbb{Z}$ -algebras provides an appropriate framework for “noncommutative projective geometry”, and Theorem 5.2 shows that the filtered $\mathbb{Z}$ -algebra $B^\lambda(\chi)$ deserves to be called a noncommutative deformation of the minimal resolution $\tilde{X} \rightarrow X$.

5.5. On the other hand, as explained in §1.3 of [GS], Theorem 5.2 naturally completes the following diagram (which does not commute!):

$$\begin{array}{ccc} ? & \xleftarrow{\sim} & \mathcal{O}^\lambda\text{-mod} \\ \text{gr} \downarrow & & \downarrow \text{gr} \\ \text{Coh}(\tilde{X}) & \xleftarrow{\text{pullback}} & \text{Coh}(X) \end{array}$$

Namely, the question mark can be replaced by the category $B^\lambda(\chi)\text{-qgr}$. The corresponding fact for rational Cherednik algebras of type A has been proved in [GS], and then applications of this result to the representation theory of these algebras have been explored in [GS2]. A similar study for the algebras $\mathcal{O}^\lambda$ will appear in [B].

6. PROOF OF THE MAIN THEOREM

6.1. In this section we prove Theorem 5.2. First we collect several unrelated facts used in the proof, some of which may be well known to the experts in the corresponding areas of mathematics, but which we explain here for the reader's convenience (and for the lack of suitable references). The proof of the theorem itself begins in §6.7.

6.2. We start with a general geometric statement.

Proposition 6.1. *Let X be a normal affine irreducible algebraic surface over $\mathbb{C}$, and $\mathcal{E}$ a torsion-free coherent sheaf on X . Then $\text{Ext}^n(\mathcal{E}, \mathcal{O}_X)$ is finite dimensional for all $n \geq 1$. In addition, if $\mathcal{E}$ has generic rank 1, then*

$$\text{End}_{\mathcal{O}_X}(\mathcal{E}) = \Gamma(X, \mathcal{O}_X). \quad (6.1)$$

Proof. Note that X has at worst finitely many singular points because it is a normal surface. As usual, we write $\mathcal{E}^\vee = \mathcal{H}om(\mathcal{E}, \mathcal{O}_X)$ for the dual sheaf. The canonical map $\mathcal{E} \rightarrow \mathcal{E}^{\vee\vee}$ is known to be an isomorphism away from the singular points of X and possibly another finite set of points; moreover, $\mathcal{E}^{\vee\vee}$ is locally free away from the singular points of X . This implies that $\mathcal{E}$ itself is also locally free away from a finite set of points. Thus for $n \geq 1$, the sheaf $\mathcal{E}xt^n(\mathcal{E}, \mathcal{O}_X)$ has finite support. On the other hand, since X is affine, the category of coherent $\mathcal{O}_X$ -modules is equivalent to the category of finitely generated $\mathbb{C}[X]$ -modules, and in particular $\text{Ext}^n(\mathcal{E}, \mathcal{O}_X) \cong \Gamma(X, \mathcal{E}xt^n(\mathcal{E}, \mathcal{O}_X))$. This implies the first statement.

For the second statement, let $S \subset X$ be the finite set of points where $\mathcal{E}$ is not locally free. Then

$$\Gamma(X, \mathcal{O}_X) \subseteq \text{End}_{\mathcal{O}_X}(\mathcal{E}) \hookrightarrow \text{End}_{\mathcal{O}_{X \setminus S}}(\mathcal{E}|_{X \setminus S}) = \Gamma(X \setminus S, \mathcal{O}_X) = \Gamma(X, \mathcal{O}_X),$$

where the second inclusion follows from the assumption that $\mathcal{E}$ is torsion-free, and the last equality follows from the assumption that X is normal. This completes the proof. $\square$

6.3. We now prove two results concerning the algebras $\mathcal{O}^\lambda$ that can be deduced from [CBH].

Proposition 6.2. *Let $\lambda \in \mathbb{C}^I$ be such that $\lambda \cdot \delta = 1$ and $\mathcal{O}^\lambda$ is regular. Given $d \in \mathbb{N}$, the algebra $\mathcal{O}^{\lambda+\xi}$ has no nonzero modules of dimension $\leq d$ for all sufficiently large $\xi \in \Lambda_+$.*

Proof. Obviously, it suffices to prove the statement above with the word “nonzero” replaced by the word “simple”. We now recall from [CBH] that the deformed preprojective algebra Π^λ defined in §1.1 is Morita equivalent to $\mathcal{O}^\lambda$, the equivalence being given explicitly by

$$M \mapsto e_0 \Pi^\lambda \otimes_{\Pi^\lambda} M = e_0 M,$$

where $e_0 \in \Pi^\lambda$ denotes the idempotent corresponding to the extending vertex. Note that this functor acts as $\alpha \mapsto e_0 \cdot \alpha$ on the dimension vectors of the modules.

Now we know from [CBH] that the dimension vectors of simple finite dimensional Π^λ -modules are among the positive roots α satisfying $\lambda \cdot \alpha = 0$ (observe that all such roots are necessarily real and non-Dynkin). Also, given $\xi \in \Lambda$, the map

$$\alpha \mapsto \alpha - (\xi \cdot \alpha)\delta$$

establishes a bijection of *finite* sets

$$\left\{ \text{roots } \alpha \text{ such that } \lambda \cdot \alpha = 0 \right\} \xrightarrow{\sim} \left\{ \text{roots } \beta \text{ such that } (\lambda + \xi) \cdot \beta = 0 \right\}.$$

If α is a real root, let us write $\alpha' = \alpha - (\epsilon_0 \cdot \alpha)\delta$, which is a Dynkin root. Observe that if $\xi \in \Lambda$, then we have $\xi \cdot \alpha = \xi \cdot \alpha'$. Thus, if $\beta = \alpha - (\xi \cdot \alpha)\delta$, then

$$\epsilon_0 \cdot \beta = \epsilon_0 \cdot \alpha - \xi \cdot \alpha'. \quad (6.2)$$

Now, using finiteness, choose $N \in \mathbb{N}$ such that $|\epsilon_0 \cdot \alpha| \leq N$ for every root α with $\lambda \cdot \alpha = 0$, and choose $\xi_0 \in \Lambda_+$ such that

$$\xi_0 \cdot \psi > N + d \quad \text{for every positive Dynkin root } \psi.$$

It follows that if $\xi \geq \xi_0$ and α, β are related as above, then (6.2) implies

$$|\epsilon_0 \cdot \beta| \geq |\xi \cdot \alpha'| - |\epsilon \cdot \alpha| > N + d - N = d.$$

Finally, note that $\mathcal{O}^{\lambda+\xi}$ is regular for sufficiently large ξ , since, according to [CBH], we only need to choose ξ large enough so that $(\lambda + \xi) \cdot \alpha \neq 0$ for every Dynkin root α . It then follows from the discussion above that the possible dimensions of the simple finite dimensional $\mathcal{O}^{\lambda+\xi}$ modules are among the integers $\epsilon_0 \cdot \beta$, where β is a positive root with $(\lambda + \xi) \cdot \beta = 0$, which completes the proof of the proposition. $\square$

Remark 6.3. We have stated and proved the proposition above for left $\mathcal{O}^\lambda$ -modules. However, the same result also holds for right modules. Indeed, if (Q, I) is any quiver, it is easy to see that for any $\lambda \in \mathbb{C}^I$ there is a natural isomorphism between $\Pi^{-\lambda}(Q)$ and the opposite algebra of $\Pi^\lambda(Q)$ that preserves the idempotents corresponding to the vertices of Q . In particular, it follows from [CBH] that the dimension vectors of simple left $\Pi^\lambda(Q)$ -modules and simple right $\Pi^\lambda(Q)$ -modules are the same.

Proposition 6.4. *Let $\lambda \in \mathbb{C}^I$ and $\xi \in \Lambda$ be such that $\lambda \cdot \delta = 1$ and the algebras $\mathcal{O}^\lambda$ and $\mathcal{O}^{\lambda+\xi}$ are both regular. Then there exists a Morita equivalence*

$$\mathcal{O}^\lambda\text{-mod} \xrightarrow{\sim} \mathcal{O}^{\lambda+\xi}\text{-mod}, \quad (6.3)$$

compatible with filtrations in the obvious sense.

Proof. Under the assumptions of the proposition, the results of [CBH] imply that the functors $\Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} -$ and $e_0 \Pi^{\lambda+\xi} \otimes_{\Pi^{\lambda+\xi}} -$ provide equivalence of categories

$$\mathcal{O}^\lambda\text{-mod} \xrightarrow{\sim} \Pi^\lambda\text{-mod} \quad \text{and} \quad \Pi^{\lambda+\xi}\text{-mod} \xrightarrow{\sim} \mathcal{O}^{\lambda+\xi}\text{-mod}$$

that are compatible with filtrations. Hence, if we can show that there is an equivalence $\Pi^\lambda\text{-mod} \xrightarrow{\sim} \Pi^{\lambda+\xi}\text{-mod}$ that is also compatible with filtrations, we can define (6.3) as the composition of these three equivalences.

Now let E denote the affine space of all $\lambda \in \mathbb{C}^I$ such that $\lambda \cdot \delta = 1$. Each simple reflection s_i corresponding to a vertex $i \in I$ of Q defines an automorphism $r_i : E \rightarrow E$, and the reflection functors of [CBH] provide an equivalence $\Pi^\lambda\text{-mod} \xrightarrow{\sim} \Pi^{r_i\lambda}\text{-mod}$ for every $\lambda \in E$. It is clear from the construction given in loc. cit. that this equivalence is compatible with filtrations. On the other hand, if $\phi : Q \rightarrow Q$ is an automorphism of the underlying graph of Q , it also induces an automorphism $\phi^* : E \rightarrow E$, and it is easy to see that there is a natural isomorphism $\Pi^\lambda \rightarrow \Pi^{\phi^*\lambda}$ of filtered algebras, for any $\lambda \in E$. Thus we have reduced the proof of the proposition to Lemma 6.5 below. $\square$

Lemma 6.5. *Given $\xi \in \Lambda$, the map $\lambda \mapsto \lambda + \xi$ can be written as a composition of simple reflections and automorphisms of the graph Q .*

Proof. We use some standard facts about root systems and affine Weyl groups that can be found in Chapter VI of [Bour]. Consider the vector space $V = (\mathbb{Z}^I/\mathbb{Z}\delta) \otimes_{\mathbb{Z}} \mathbb{C}$; it is well known that the image of the set of real roots for Q under the projection map $\mathbb{Z}^I \rightarrow \mathbb{Z}^I/\mathbb{Z}\delta$ is a reduced root system $\mathcal{R}$ in the space V , in the sense of loc. cit., Chapter VI, §1.4. Now V^* is naturally identified with $\Lambda \otimes_{\mathbb{Z}} \mathbb{C}$, and E can be viewed as an affine space for the vector space V^* . Let W_{ext} denote the group of automorphisms of E generated by the translations by the elements of Λ and by the Weyl group W_{fin} of the root system $\mathcal{R}$; sometimes W_{ext} is called the *extended Weyl group* of the root system $\mathcal{R}$. It follows from the results of loc. cit., Chapter VI, §2.1 and §2.3, that W_{ext} has the alternate description as the group of automorphisms of E generated by the *affine* Weyl group W_{aff} (which by definition is generated by the simple reflections corresponding to all vertices of Q ; it is called simply the Weyl group of Q in [CBH]) and the group of automorphisms of the graph Q . This proves the lemma. $\square$

Remark 6.6. As we will see below, our proof of Theorem 5.2 relies heavily on the fact that the “shift functors”

$$P_\xi^\lambda \otimes_{\mathcal{O}^\lambda} - : \mathcal{O}^\lambda\text{-mod} \longrightarrow \mathcal{O}^{\lambda+\xi}\text{-mod}, \quad (6.4)$$

where the bimodules P_ξ^λ have been defined in §5.3, are equivalences of categories for “sufficiently large” λ and ξ . Even though the proof of Proposition 6.4 provides a definition of shift functors, it seems impractical to use this result for quantization of minimal resolutions of Kleinian singularities, since it is hard to compute explicitly the associated graded spaces of the bimodules defining the equivalences (6.3), and to control the compositions of these equivalences. It is not known to us if the shift functors of Proposition 6.4 are isomorphic to the shift functors (6.4).

6.4. Next we prove a characterization of Morita equivalence which is somewhat different from the ones found in standard textbooks, and for which we don’t know a reference.

Proposition 6.7. *Let A, B be rings, and let P be an (A, B) -bimodule such that the natural ring homomorphisms*

$$A \longrightarrow \text{End}_{\text{Mod-}B}(P) \quad \text{and} \quad B \longrightarrow \text{End}_{A\text{-Mod}}(P)^{op}$$

are isomorphisms. If P is projective both as a left A -module and as a right B -module, then the functor $P \otimes_B -$ gives a Morita equivalence between B and A .

Proof. We freely use the terminology and results of §3.5 of [MCR]. In particular, if M is a left module over a ring R , and $S = \text{End}_{R\text{-Mod}}(M)^{\text{op}}$, then in order to show that M yields a Morita equivalence between R and S , it is enough to prove that M is a projective R -module, and that it is a generator of the category $R\text{-Mod}$.

Let us write

$$P^* = \text{Hom}_{A\text{-Mod}}(P, A) \quad \text{and} \quad P^\vee = \text{Hom}_{\text{Mod-}B}(P, B);$$

both spaces have the natural (B, A) -bimodule structures. In view of the previous paragraph, it suffices to show that P is a generator of the category $A\text{-Mod}$, in other words, that the evaluation map

$$\text{ev} : P \otimes_B P^* \longrightarrow A \tag{6.5}$$

is surjective. The fact that P is a projective right B -module implies that the natural map

$$\phi : P \otimes_B P^\vee \longrightarrow \text{End}_{\text{Mod-}B}(P) \xleftarrow{\simeq} A$$

given by sending $x \otimes f$ to the endomorphism $y \mapsto x \cdot f(y)$ is surjective. However, it is clear that this map must factor through the map (6.5). More precisely, if $\psi : P^\vee \rightarrow P^*$ is defined by $\psi(f)(x) = \phi(x \otimes f)$, then $\phi = \text{ev} \circ (1 \circ \psi)$. This shows that (6.5) is surjective, completing the proof of the proposition. $\square$

6.5. In this subsection we recall, following §2.2 of [MCR], the notion of a uniform module, and explain how it will be applied in our proof of Theorem 5.2. Let A be a ring and M a (left) A -module. We say that a submodule $N \subseteq M$ is *essential* if $N \cap X \neq (0)$ for every nonzero submodule $X \subseteq M$. We say that M is *uniform* if $M \neq (0)$ and every nonzero submodule of M is essential.

Lemma 6.8. *Let B be a left Ore domain and P an (A, B) -bimodule which yields a Morita equivalence between A and B . Then P is a uniform A -module. In particular, if P' is any torsion-free nonzero left A -module, then every surjective homomorphism $f : P \rightarrow P'$ of A -modules must be an isomorphism.*

Proof. Note that the equivalence $B\text{-mod} \xrightarrow{\sim} A\text{-mod}$ induced by P maps B to P . Also, B is a uniform left B -module by the last remark of §2.2.5 of loc. cit. This implies that P is a uniform left A -module, by Lemma 3.5.8(vi) of loc. cit. In particular, if $f : P \rightarrow P'$ is as in the statement of the lemma, and $K = \text{Ker}(f) \neq (0)$, then K is an essential submodule of P . If $x \in P$ is any element such that $f(x) \neq 0$, then we see that $ax \in K$ for some $a \in A$, which implies that $a \cdot f(x) = 0$, contradicting the assumption that P' is torsion-free. $\square$

6.6. We now record some observations in noncommutative ring theory that are interesting in their own right. For the definition of and basic facts about rigid dualizing complexes, as well as the notions of Cohen-Macaulay rings and Gorenstein rings in the noncommutative situation, we refer the reader to §3 of [EG] and references therein.

Proposition 6.9. *Let A be a finitely generated connected filtered algebra such that $\text{gr}^\bullet A$ is commutative and Gorenstein.*

- (1) If M is a finitely generated left A -module equipped with a good filtration, and $\mathrm{gr}^\bullet M$ is a Cohen-Macaulay $\mathrm{gr}^\bullet A$ -module of degree k , then M is a Cohen-Macaulay A -module of degree k .
- (2) If, in addition, A is regular and M is a finitely generated Cohen-Macaulay A -module of degree 0, then M is projective.

Proof. The argument for (1) is essentially given in §3 of [EG]. Namely, since $\mathrm{gr}^\bullet A$ is commutative and Gorenstein, it is its own dualizing complex, and then the same is true of A . But for every $j \geq 0$, a result of Björk ([Bj], Proposition 3.1) shows that $\mathrm{gr}^\bullet \mathrm{Ext}_A^j(M, A)$ is a subquotient of $\mathrm{Ext}_{\mathrm{gr}^\bullet A}^j(\mathrm{gr}^\bullet M, \mathrm{gr}^\bullet A)$. Thus if $\mathrm{Ext}_{\mathrm{gr}^\bullet A}^j(\mathrm{gr}^\bullet M, \mathrm{gr}^\bullet A) = (0)$ for all $j \neq k$, then $\mathrm{Ext}_A^j(M, A) = (0)$ for all $j \neq k$, which proves (1).

For (2), in view of the previous paragraph, it suffices to show that if $\mathrm{Ext}_A^j(M, A) = (0)$ for all $j \geq 1$, then M is projective. We use induction on the projective dimension of M , which is finite by assumption. If the projective dimension is 0, we are done. Otherwise there is an exact sequence

$$0 \longrightarrow N \longrightarrow P \longrightarrow M \longrightarrow 0 \quad (6.6)$$

with P finitely generated and projective. The projective dimension of N is one less than that of M , and the long exact sequence of Ext 's shows that $\mathrm{Ext}_A^j(N, A) = (0)$ for all $j \geq 1$, whence N is projective by induction. In particular, our assumption on M implies that $\mathrm{Ext}_A^1(M, N) = (0)$, so the sequence (6.6) splits, and thus M is projective. $\square$

6.7. We return to the situation of Theorem 5.2. Given the construction of the minimal resolution $\tilde{X} \rightarrow X$ discussed in Section 2, it is clear that we only need to prove the first part of the theorem, namely, that the $\mathbb{Z}$ -algebra $B(\lambda + \xi, \chi)$ satisfies

$$\mathcal{O}^\lambda\text{-mod} \simeq B(\lambda + \xi, \chi)\text{-qgr}$$

for all sufficiently large ξ . Using Proposition 6.4 and Theorem 4.4, we see that it is enough to check that $B(\lambda + \xi, \chi)$ is a Morita $\mathbb{Z}$ -algebra, in the sense of Definition 4.3, for all sufficiently large ξ . We combine the results of Section 2 and §§6.3–6.6. The second condition in the definition of what it means for $B(\lambda + \xi, \chi)$ to be a Morita $\mathbb{Z}$ -algebra follows from the first one in view of Theorem 2.2(3) and Lemma 6.8. Thus we need to check that there exists a large enough $\xi \in \Lambda_{++}$ for which the $(\mathcal{O}^{\lambda+\xi+i\chi}, \mathcal{O}^{\lambda+\xi+j\chi})$ -bimodules

$$P_{ij} := P_{(i-j)\chi}^{\lambda+\xi+j\chi}$$

satisfy the assumptions of Proposition 6.7 whenever $i - j \gg 0$.

6.8. Let $N \in \mathbb{N}$ be fixed, so that the statements of the last two parts of Theorem 2.2 hold for all $m, n \geq N$. The key point is that, *whereas the bimodules P_{ij} themselves depend on ξ , the associated graded modules do not*. In fact, Theorem 3.2 implies that $\mathrm{gr}^\bullet P_{ij} \cong S_{i-j}$ as R -bimodules, using the notation of Section 2. Now the second statement of Proposition 6.1

and the standard argument given, e.g., in §3 of [EG], in the proof of Theorem 1.5(iv), implies that for $i - j \geq N$, we have

$$\mathcal{O}^{\lambda+\xi+i\chi} \xrightarrow{\simeq} \text{End}_{\text{mod-}\mathcal{O}^{\lambda+\xi+j\chi}}(P_{ij})$$

and

$$\mathcal{O}^{\lambda+\xi+j\chi} \xrightarrow{\simeq} \text{End}_{\mathcal{O}^{\lambda+\xi+i\chi}\text{-mod}}(P_{ij})^{op}.$$

Hence we only need to make sure that P_{ij} is projective as a left and as a right module.

For simplicity, let us write $\mathcal{O}_i = \mathcal{O}^{\lambda+\xi+i\chi}$. Now for $n = N, N+1, \dots, 2N-1$, the first statement of Proposition 6.1 and the argument used in the proof of Proposition 6.9(1) imply that if $i - j = n$, then the modules

$$\text{Ext}_{\mathcal{O}_i\text{-mod}}^\ell(P_{ij}, \mathcal{O}_i) \quad \text{and} \quad \text{Ext}_{\text{mod-}\mathcal{O}_j}^\ell(P_{ij}, \mathcal{O}_j) \quad (\ell = 1, 2) \quad (6.7)$$

have finite dimension which is *independent of* ξ , and depends only on $i - j$ but not on i or j separately. Furthermore,

$$\text{Ext}_{\mathcal{O}_i\text{-mod}}^\ell(P_{ij}, \mathcal{O}_i) = \text{Ext}_{\text{mod-}\mathcal{O}_j}^\ell(P_{ij}, \mathcal{O}_j) = (0) \quad \text{for } \ell \geq 3,$$

because the global dimensions of $\mathcal{O}_i$ and $\mathcal{O}_j$ are at most 2 (see [CBH]). In particular, by Proposition 6.2 and Remark 6.3, we can choose and fix a large enough ξ for which the modules (6.7) are necessarily zero if $i - j \in \{N, N+1, \dots, 2N-1\}$. Thus, by Proposition 6.9(2), we have now shown that, for the ξ that we've chosen, the bimodules P_{ij} induce Morita equivalences between the algebras $\mathcal{O}_j$ and $\mathcal{O}_i$ whenever $i - j \in \{N, N+1, \dots, 2N-1\}$.

Remark 6.10. Note that if $\mathcal{O}^\lambda$ has no nonzero finite dimensional modules (by Theorem 0.3 of [CBH], this happens if and only if $\lambda \cdot \alpha \neq 0$ for all non-Dynkin roots α), then the modules (6.7) are automatically zero, and the choice of ξ is unnecessary. We conjecture that, in fact, the modules P_{ij} induce Morita equivalences between the algebras $\mathcal{O}_i$ and $\mathcal{O}_j$ provided λ is *dominant* in the sense that $\text{Re}(\lambda \cdot \alpha) > 0$ for every positive Dynkin root α .

6.9. We complete the proof as follows. With the notation above, it is obvious that any integer $m \geq 2N-1$ can be written as a sum $m = m_1 + \dots + m_k$, where each $m_s \in \{N, N+1, \dots, 2N-1\}$. Thus, for $i - j = m$, we see from Theorem 2.2(3) that the bimodule P_{ij} is a homomorphic image of the tensor product

$$\mathcal{P} := P_{j+m_1, j} \otimes P_{j+m_2, j+m_1} \otimes \dots \otimes P_{i, j+m_{k-1}}$$

over the appropriate algebras $\mathcal{O}_{j+m_s}$. Since each factor in the tensor product induces a Morita equivalence by the argument above, so does the tensor product. Hence, by Lemma 6.8, the surjection $\mathcal{P} \rightarrow P_{ij}$ must necessarily be an isomorphism; in particular, P_{ij} also induces a Morita equivalence between $\mathcal{O}_j$ and $\mathcal{O}_i$, and the proof of the theorem is complete.

REFERENCES

- [Bj] J.-E. Björk, *The Auslander condition on Noetherian rings*, Sémin. d'Algèbre Paul Dubreil et Marie-Paul Malliavin, 39ème Année (Paris, 1987/1988), 137–173, Lect. Notes in Math. **1404**, Springer, Berlin, 1989.
- [Bour] N. Bourbaki, “Éléments de mathématique, fasc. 34: Groupes et algèbres de Lie, Chapitres IV–VI.” Actualités Sci. Indust. **1337**, Hermann, Paris, 1968.
- [B] M. Boyarchenko, *Quantization of minimal resolutions of Kleinian singularities II: complements and applications*, in preparation.
- [CS] H. Cassens and P. Slodowy, *On Kleinian singularities and quivers*, in: “Singularities (Oberwolfach, 1996),” 263–288, Progr. Math. **162**, Birkhäuser, Basel, 1998.
- [C-B] W. Crawley-Boevey, *Geometry of the moment map for representations of quivers*, Compositio Math. **126** (2001), no. 3, 257–293.
- [CBH] W. Crawley-Boevey and M. Holland, *Noncommutative deformations of Kleinian singularities*, Duke Math. J. **92** (1998), no. 3, 605–635.
- [EG] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, Invent. Math. **147** (2002), no. 2, 243–348.
- [GS] I. Gordon and J.T. Stafford, *Rational Cherednik algebras and Hilbert schemes*, Preprint, July 2004, [math.RA/0407516](#)
- [GS2] I. Gordon and J.T. Stafford, *Rational Cherednik algebras and Hilbert schemes II: representations and sheaves*, Preprint, October 2004, [math.RA/0410293](#)
- [EGA II] A. Grothendieck, “Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes.” Inst. Hautes Études Sci. Publ. Math. **8** (1961).
- [Ho] M.P. Holland, *Quantization of the Marsden-Weinstein reduction for extended Dynkin quivers*, Ann. Sci. École Norm. Sup. (4) **32** (1999), no. 6, 813–834.
- [MCR] J.C. McConnell and J.C. Robson, “Noncommutative Noetherian Rings.” Grad. Studies in Math. **30**, AMS, Providence, RI, 2001.
- [GIT] D. Mumford, J. Fogarty and F. Kirwan, “Geometric Invariant Theory,” 3rd ed. Ergeb. Math. Grenzgeb. (2) **34**. Springer-Verlag, Berlin, 1994.
- [Mu] I.M. Musson, *Noncommutative deformations of type A Kleinian singularities and Hilbert schemes*, Preprint, April 2005, [math.RT/0504543](#)